

Radiative electron attachment to molecules of astrophysical interest: Direct and indirect mechanisms


V. Kokoouline¹, M. Khamesian¹, N. Douguet², S. Fonseca dos Santos², O. Dulieu³, M. Raoult³, A. E. Orel²

¹Department of Physics, University of Central Florida, Orlando, FL


²Department of CHMS, University of California at Davis, Davis, CA 95616, USA

³Laboratoire Aimé Cotton, CNRS, Université de Paris Sud, Orsay, France

A first-principle theoretical approach to study the process of radiative electron attachment is developed and applied to the negative molecular ions CN^- , C_4H^- , and C_2H^- . Among these anions, the first two have already been observed in the interstellar space. Cross sections and rate coefficients for formation of these ions by radiative electron attachment to the corresponding neutral radicals are calculated. For completeness of the theoretical approach, two pathways for the process have been considered: (i) A direct pathway, in which the electron in collision with the molecule spontaneously emits a photon and forms a negative ion in one of the lowest vibrational levels, and (ii) an indirect, or two-step pathway, in which the electron is initially captured through non-Born-Oppenheimer coupling into a vibrationally resonant excited state of the anion, which then stabilizes by radiative decay. We have developed a general theoretical model to describe the two pathways. The contribution of the indirect pathway to the formation of cosmic anions was found to be negligible in comparison to the direct mechanism. The obtained total rate coefficients of radiative electronic attachment at 30K are $7 \times 10^{-16} \text{ cm}^3/\text{s}$ for CN^- , $7 \times 10^{-17} \text{ cm}^3/\text{s}$ for C_2H^- , and $2 \times 10^{-16} \text{ cm}^3/\text{s}$ for C_4H^- . These rates weakly depend on temperature between 10K and 100 K. The validity of our calculations is verified by comparing the present theoretical results with data from recent photodetachment experiments performed for the CN^- , C_4H^- , and C_2H^- ions.



Calculated rate coefficients for radiative electron attachment to the CN^- , C_2H^- , and C_4H^- molecules.



Comparison of the present theoretical cross sections with available experimental data for the process of photodetachment of the CN^- , C_2H^- , and C_4H^- ions.

This work is supported by the DOE Office of Basic Energy Science and the National Science Foundation, Grant No's PHY-08-55092 and PHY-08-55622.